

Pathogen Management Plan

2013 - 2016

joondalup.wa.gov.au



Contents

Section Page 1.0 Introduction 3 1.1 Purpose of the Plan 4 1.2 Strategic Context 5 1.3 What are Pathogens? 6 7 1.4 Phytophthora 1.5 Armillaria 10 2.0 Pathogen Risk Analysis 12 2.1 Introduction 12 2.2 Study Area 12 14 2.3 Pathogen Assessment and Mapping 2.4 High Resolution Multi-Spectral Imagery 16 3.0 Pathogen Management 17 3.1 Pathogen Interpretation and Mapping 18 3.2 Bushland Management 18 3.3 Hygiene Measures 18 3.4 Access 19 3.5 Signage 19 3.6 Communication and Education 20 3.7 Partnerships 21 3.8 Treatment 21 22 3.9 Monitoring and Reporting 4.0 Management Plan Review 23 5.0 References 24 6.0 Appendices 25

Acknowledgements

Mr Paul Barber, Arbor Carbon, Environmental and Arboricultural Consultants Mr Mike Stukely, Vegetation Health Services – Science Division, Department of Environment and Conservation

Professor Giles Hardy, Centre for *Phytophthora* Science and Management, Murdoch University



1. Introduction

The City of Joondalup is situated along the Swan Coastal Plain, 30 kilometres from the Perth Central Business District. The City covers an area of 96.5 kilometres which encompasses a diverse range of natural areas including 17 kilometres of coastal foreshore, a chain of wetlands and a variety of bushland ecosystems.

The City's southern boundary is located approximately 16 kilometres from the Perth Central Business District, and is bounded by the City of Wanneroo to the east and north, the City of Stirling to the south, and the Indian Ocean to the west.

There are a variety of regionally, nationally and internationally significant natural areas located within the City including the Yellagonga Regional Park, the Marmion Marine Park, the Neerabup National Park and a number of Bush Forever sites which contain species of high conservation value.

Vegetated areas within the City are at risk from pathogens which pose a serious threat to the biodiversity values within the City's parks and bushland areas. Effective pathogen management is required to ensure that measures are taken to mitigate the effects and limit the spread of pathogens within the City.

In order to protect native vegetation and ecosystems within the City of Joondalup a Pathogen Management Plan has been developed. The Plan identifies the level of risk for pathogens becoming introduced to City parks and natural areas and prioritises the areas for further on ground investigations.

The Plan also provides guidance on the management of pathogens within the City in order to minimise the spread of pathogens. Strategies to engage the community in order to raise the awareness of pathogens within the City of Joondalup are also identified within the Plan.



Shepherds Bush Conservation Area Kingsley.

1.1 Purpose of the Plan

The Pathogen Management Plan aims to establish the level of risk of pathogens within City parks and natural areas and identifies areas of high risk where management actions should be concentrated.

The Pathogen Management Plan includes the following:

- Desktop assessment of parks and natural areas for pathogens including *Phytophthora* species and *Armillaria* to establish the level of risk within the City;
- Development of preventive and management strategies and procedures to be employed within activities relating to the use of City parks, streetscapes and natural areas;
- Identification of control and treatment measures for infested areas; and
- Development of education and communication mechanisms to raise the awareness of pathogens within the organisation and the community.

The objective of the Pathogen Management Plan is to protect biodiversity values within the City of Joondalup by minimising the risk of introducing and spreading pathogens including species of *Phytophthora* and *Armillaria luteobubalina* within landscaped and natural areas of the City.



1.2 Strategic Context

The purpose of the Pathogen Management Plan aligns with the environmental aims and objectives of a number of City of Joondalup Plans including:

Strategic Plan

The *City of Joondalup Strategic Plan 2012 – 2022* highlights the focus on preservation, rehabilitation and maintenance of the City's natural environmental assets and the importance of engaging with the community, key stakeholders and relevant agencies.


Environment Plan

The *City of Joondalup Environment Plan 2007 – 2011* identifies the key environmental pressures and threats and provides the strategic response to the major issues affecting the City of Joondalup.

Biodiversity Action Plan

The City of Joondalup *Biodiversity Action Plan 2009 – 2019* provides direction for the City's biodiversity management activities and recommends the development of Dieback Management Plans as an action.

Figure 1- City of Joondalup Strategic Environmental Framework



1.3 What are Pathogens?


Organisms such as fungi, bacteria and viruses that cause plant diseases are known as pathogens. Whilst some pathogens are naturally occurring within soil populations, others have been introduced to the environment through the movement of plant materials and soils.

Pathogens can exist in the soil for long periods of time without causing an outbreak of disease in plants. Disease outbreaks are either caused by an increase in the population of the pathogen, by an increase in the susceptibility of the plant or a change in environmental conditions. The population of the pathogen is dependent on whether the soil conditions are favourable for its growth and survival. The conditions that are favourable for the growth and survival of pathogens are different for each species of pathogen but are related to:

- Soil pH;
- Soil water content;
- Soil oxygen level;
- Nutrient levels in soil; and
- The activities of other soil organisms.

The symptoms produced by plants that are affected by pathogens vary depending upon the species of pathogen, host species, environment and climatic conditions. Some pathogens can cause rapid death of plants whilst others result in a slow, perennial decline in health.

The City of Joondalup Pathogen Management Plan provides information and management recommendations to address species of the pathogens *Phytophthora* and *Armillaria*.



1.4 Phytophthora

Ecology and Biology

The scientific name *Phytophthora* (pronounced fy-toff-thor-ah) is derived from the Greek words meaning 'plant destroyer'. *Phytophthora* is a water mould which attacks the roots and stem tissue of living plants causing them to rot. The rotting weakens and kills the plant by limiting the uptake of water and nutrients, as detailed in *Figure 2*.




Figure 2- Lifecycle of Typical Phytophthora (Sourced from Dieback Working Group).

Phytophthora cinnamomi

Phytophthora cinnamomi, also known as Phytophthora Dieback, is one of the biggest threats to biodiversity within Australia. The Environment Protection and Biodiversity Conservation Act 1999 lists 'Dieback caused by the root-rot fungus Phytophthora cinnamomi as a key threatening process to Australia's biodiversity'.

Phytophthora cinnamomi, occurs widely across southern Australia and is most commonly found in areas which receive above average of 600mm of annual rainfall, however the pathogen has been known to extend into areas which receive less rainfall. *Phytophthora cinnamomi* is found in Tasmania, Victoria, New South Wales, Queensland, South Australia and Western Australia.

Phytophthora Dieback is causing significant damage in the South-West Australia Ecoregion of Western Australia, as illustrated in *Figure 3*, because:

- Over 40% of native plant species and over 50% of rare or endangered flora species in the region are susceptible;
- The climate and soils of the South-West Australia Ecoregion suit the pathogen's survival and spread; and

• The pathogen was spread widely before it was identified as the cause of permanent damage to our ecosystems.

Thought to be an introduced pathogen, possibly originating in South-East Asia, *Phytophthora cinnamomi* is widespread throughout the urban areas of Perth. This pathogen has a very wide host range, with more than 2300 species of Western Australian native plants considered to be susceptible. *Phytophthora cinnamomi* can survive for long periods in unfavourable conditions and prefers nutrient poor soils and is not considered to be as damaging within soils with a high pH soils such as those within the Quindalup complex (found along coastal areas).

Phytophthora cinnamomi can cause rapid death of plants or cause a slow, perennial decline in health of the crown and is commonly associated with the movement of dieback fronts throughout bushland, killing large numbers of highly susceptible species, earning it the name of the 'biological bulldozer'.

Whilst *Phytophthora cinnamomi* is the most common species within Western Australia other species of the pathogen have also been detected within Western Australia and the Perth Metropolitan Region, these species include:

- Phytophthora multivora
- Phytophthora humicolalike
- Phytophthora nicotianae.

Phytophthora multivora

Phytophthora multivora is widespread throughout the South-West of Western Australia with a similar distribution to *Phytophthora cinnamomi*. However unlike *Phytophthora cinnamomi*, it seems to tolerate the higher pH soils of the Quindalup and Spearwood dune systems, and is now one of the most commonly encountered *Phytophthora* species associated with disease symptoms of native trees on these soil types.



Figure 3- Distribution of Phytophthora within the South-West region of Western Australia.

there is no data on the pathogenicity (ability to produce disease in a host organism) of *Phytophthora humicolalike* to hosts plants that reside within the Perth Metropolitan Region.

The known distribution of *Phytophthora humicolalike* in Western Australia is limited to a small number of cases in Perth and Esperance. In these cases the pathogen was isolated either directly from streams, or from a site periodically inundated with water, suggesting the species may be well adapted to surviving in such conditions, meaning that wetland ecosystems may be at the greatest risk from this species of *Phytophthora*.

Very little is known about the host range of *Phytophthora humicolalike*, other than it being identified in *Casuarina obesa*. Like other *Phytophthora*, species, it is suspected to be transmitted to other trees via root to root contact and through contaminated soil or plant material.

As *Phytophthora humicolalike* is a newly identified species of *Phytophthora,* research into the susceptibility and resistance of host plants has not yet been undertaken.

Suspected Phytophthora multivora within a City of Joondalup Park.

Phytophthora multivora is common in urban areas of Perth, particularly along the inland dune systems. Probability of entry and establishment may also be higher in parks where containerized nursery stock and soil is introduced, and the use of machinery and vehicles is common.

The symptoms produced on plants vary depending upon the host, environment and climatic conditions; however Phytophthora multivora can cause rapid death of plants, or a slow, perennial decline in health of the crown. It has been observed causing large, aggressive lesions on Banksia species but on other species such as Tuarts (Eucalyptus gomphocephala), is considered to be a pathogen of the fine roots only. Unlike Phytophthora *cinnamomi*, it is not associated with 'dieback fronts' but is more commonly associated with individual spot deaths and areas of tree decline.

Phytophthora humicolalike

Phytophthora humicolalike is a new species of Phytophthora yet to be formally described. Biologically it is similar in characteristics to other species of Phytophthora. Phytophthora humicolalike has been associated with decline and death of Casuarina species in parts of Western Australia. Currently



Phytophthora nicotianae

Phytophthora nicotianae is a soil-borne pathogen which was identified as affecting herbaceous and woody plants used within agriculture and horticulture; however it is now established within natural ecosystems of Western Australia.

Phytophthora nicotianae has been recorded from numerous host families in Western Australia including a range of native hosts. The pathogen is found in soil or plant tissue and prefers moist conditions for survival, however it can also survive for long periods in unfavourable conditions.

Phytophthora nicotianae is widely found within nursery stock and therefore has a higher probability of infecting parks and reserves than for species such as humicolalike, which has never been confirmed within nursery stock. The probability of entry and establishment in parks compared to natural areas is also higher as nursery stock and soil is regularly introduced, and the use of machinery and vehicles is common.

The symptoms produced on plants vary depending upon the host, environment and climatic conditions. It has been associated with large lesions at the base of *Eucalyptus* trees and causes collar rot of *Grevillea* species. *Phytophthora nicotianae* has also been identified as causing fine root death of numerous other native plant species.

Phytophthora nicotianae can move throughout the environment via soil or water or via root to root contact between plants. This pathogen is capable of causing decline and death of seedlings, shrubs and trees and is associated with disease of vegetation throughout the Perth Metropolitan Region.

Distribution and Dispersal

A number of species of *Phytophthora* are common throughout the whole of South-West of Western Australia, including within the Perth Metropolitan Region. *Phytophthora* is easily dispersed through the movement of infected soil and plant material by humans and animals. The pathogen is commonly spread by transplanting infected plants from one site to another or from infected soil being moved on boots, tools and machinery. The application of untreated or raw green mulch is also considered a source of the pathogen.

Recreational activities within infected areas including cycling and bush walking can also lead to increased infestation. Local Government has an important role to play in managing *Phytophthora* dieback as activities such as road and drain construction and bushland management, have the potential to introduce *Phytophthora* dieback to a previously uninfected area, or increase its rate of spread.

When *Phytophthora* spreads to bushland, it kills many susceptible plants, resulting in a permanent decline in the diversity of the bushland. It can also change the composition of the bushland. Native animals that rely on susceptible plants for survival are also at risk of population decline within sites infested by *Phytophthora*.³

Different species of *Phytophthora* affect a variety of host plants with many Western Australian native plant species being susceptible to infection by several species of *Phytophthora* as detailed in Table 1.

Proteaceae	Myrtaceae	Epacridaceae	Other
Adenanthos	Agonis	Andersonia*	Allocasuarina
Banksia*	Beaufortia	Astroloma*	Anarthia
Conospermum	Calothamnus	Leucopogon*	Boronia
Dryandra	Calytrix	Lysinema*	Conostylis
Franklandia	Eremaea	Monotoca*	Dampiera
Grevillea	Eucalyptus	Sphenotoma*	Dasypogon
Hakea	Hypocalymma	Styphelia*	Daviesia
lsopogon*	Kunzea		Eutaxia
Lambertia*	Melaleuca		Gastrolobium
Persoonia*	Regelia		Hibbertia*
Petrophile*	Scholtzia		Hovea
Stirlingia*	Thryptomene*		Jacksonia
Synaphea	Verticordia*		Lasiopetalum*
Xylomelum			Latrobea
			Macrozamia
			Oxylobium
			Patersonia
			Phlebocarya
			Xanthorrhoea
			Xanthosia

* Many species in the genus are severely affected by *Phytophthora*

Table 1- Plant genera with species known to be affected by Phytophthora species (Sourced from Dieback Working Group)

1.5 Armillaria

Ecology and Biology

Armillaria luteobubalina is a soil-borne fungus that causes root rot of a wide variety of plants including many species of native flora. The fungus is native to Australia and can cause major damage to natural ecosystems. Armillaria lives and feeds on the wood of infected plants and spreads on infected roots as branching threads which can also be found under the bark of trees on the lower portion of the trunk. Armillaria reduces the function of the roots and affects the internal structure of the tree, often resulting in a slow decline in health and eventually death in trees.

Armillaria is commonly known as 'Honey Fungus' due to the colour of the fruiting bodies which grow above ground level near infected trees at certain times of the year, as illustrated in Figure 4.

Unlike some other root rot fungi, *Armillaria luteobubalina* does not infest soil. It spreads from a woody food base, which may be a tree or a stump, or even a small piece of infected root. As the root systems of many trees in these affected areas are in contact with each other, the fungus can move from its food base into the roots of a healthy tree.⁴

The pathogen commonly infects roots, root collars, basal stems and main stems of susceptible species. If infected trees are removed but the stumps remain in situ, the pathogen will rapidly colonise the stump and remaining roots, becoming a source of infection for many years.⁵

The spread and development of *Armillaria luteobubalina* is favoured by disturbance and irrigation, particularly through summer, and occurs throughout managed areas of the Perth Metropolitan Region.

The symptoms produced on plants vary depending upon the host, environment and climatic conditions. *Armillaria luteobubalina* can cause large inverted V shaped lesions at the base of trees. The fruiting bodies, seen generally in May to July in Perth, are a secondary indicator of the *Armillaria* fungus as the pathogen is much more active below ground and is usually a sign that the fungus is well established within the area.

Distribution and Dispersal

Armillaria inhabits temperate regions of Australia and can be found in a range of environments such as coastal dunes and bushland areas including those that occur along the Swan Coastal Plain.⁶

Armillaria luteobubalina has been recorded from more than 50 different plant families throughout Australia, and more than 200 species are affected, not only in native forests, woodlands and heathlands, but also in reserves, parklands and residential gardens. The pathogen commonly affects species of Eucalyptus including Jarrah and Marri. Some Banksia species are also susceptible to the pathogen.⁷

- 4. Forest Science Centre (2003)
- 6. Smith-White and Summerell, 2003
- 5. P Barber, 2012
- 7. Shearer, B. L and Tippett J. T,

Figure 4- Fruiting Bodies of Armillaria luteobubalina (Sourced from Dieback Working Group)






The pathogen does not spread via swimming spores like *Phytophthora* species. *Armillaria luteobubalina* can be spread via spores; however, it is most commonly spread via root to root contact and within diseased plant material. Movement of soil is not considered a common means of dispersal as the pathogen usually requires plant material to survive, particularly in drier sites. Untreated and green mulch is known to been a source, therefore only composted mulch should be used within landscaping activities. Transplanting infected plants from one site to another may spread the pathogen.⁸

Managing the spread of the pathogen once it has entered a site is challenging and costly. Eradication is possible by the complete removal of infected stumps and roots. Deep trenching around infected trees may prevent the spread into adjacent, uninfected areas. There are no commercial fungicides at present that can be successfully prescribed for controlling the pathogen.

Where trees and shrubs are infected and complete removal of the infected stump and lateral roots is not feasible, these trees should be retained rather than felled to minimise the colonisation of the unaffected portions of the specimen and subsequent rapid spread and infection of neighbouring vegetation. They should however be considered for removal if they are deemed to be structurally unsound. The removal of turf and grass and therefore the need for irrigation around trees during summer may also reduce the spread and development of *Armillaria luteobubalina*.

8. P Barber, 2012



2.0 Pathogen Risk Analysis

2.1 Introduction

The City of Joondalup engaged consultants Arbor Carbon to undertake a Pathogen Desktop Risk Analysis of vegetated areas within the City. The aim of the Study was to establish the level of risk for infestation of pathogen species within City parks and reserves.

The Pathogen Risk Assessment Study included:

- Identification of areas within the City at risk of pathogen infestation including the likelihood of introduction to or establishment / spread of pathogens within the area.
- Identification of vegetation communities of high priority and assessment of the level of risk pathogens pose to the area.
- Assessment of the manageability of the risk within the Study Area.
- Prioritised list of reserves for further investigations.

For the Risk Analysis consultants Arbor Carbon categorised the four species of Phytophthora and Armillaria luteobubalina under one group known as 'threatening pathogens'. Results from the Pathogen Desktop Risk Analysis Report have been utilised to inform the development of the City of Joondalup Pathogen Management Plan.

2.2 Study Area

The Study Area for the Pathogen Desktop Risk Analysis includes City parks and natural areas as illustrated in Figure 4. A list of the sites included within the Pathogen Risk Assessment Study is included within Appendix 1.





Figure 5 - Study Area - City of Joondalup Pathogen Management Plan

2.3 Pathogen Assessment and Mapping

Methodology

In order to assess the level of risk for infestation by threatening pathogens, factors affecting the likelihood for infestation were assessed against each separate park and natural area within the Study Area.

Variables that affect the level of risk which were included within the Desktop Risk Analysis include whether disease is suspected at the site (based on previous surveys), if the area is irrigated or non-irrigated, proximity to other vegetated areas, vegetation type and the level of protection and priority ranking for natural areas, derived from the Western Australian Local Government Association's Natural Areas Initial Assessment (NAIA).

Each variable was assigned a value which contributed to the overall score of the park or natural area, which was then classified as being High, Medium or Low Risk of being affected by the threatening pathogens. The overall risk rating for the area will be utilised to develop a program for on ground sampling and investigations to determine the actual extent of pathogens within City parks and reserves.

Details of the criteria and associated values used to assess the level of risk for pathogens within parks and a natural area is provided in Appendix 2.

Results

The Pathogen Desktop Risk Analysis demonstrated that there were a total of 58 parks and natural areas which are classified as High Risk (Priority 1), 210 parks and natural areas were classified as Medium Risk (Priority 2), and 112 parks were listed as Low Risk (Priority 3), as illustrated in Figure 5 and Figure 6.

Parks and natural areas within the High Risk (Priority 1) category should be given priority in regard to undertaking further on ground investigations and the implementation of pathogen management actions as these areas are at most risk of being affected by *Phytophthora* or/and *Armillaria* species.

A full list of the parks and natural areas for each of the Priority Areas for Phase 2 (Pathogen Mapping and Monitoring) is provided in Appendix 3. The Priority List and the associated values for each reserve will be utilised to inform the development of a Pathogen Interpretation and Mapping Project to establish the extent of pathogen infestation within the City.

Limitations of Desktop Study

It should be noted that the Pathogen Desktop Risk Analysis is based on the assessment of variables for the City's parks and natural areas using data sets provided by the City which includes profiles for individual vegetated areas as well as spatial information. As no on site investigations or ground truthing has been undertaken the Disease Risk for individual areas should only be used as a basis for prioritising areas for further investigations in order to establish the actual extent of threatening pathogens within individual parks and natural areas and for the prioritisation of the implementation of management actions.

The introduction of any species of *Phytophthora* or *Armillaria* into one or more City reserves poses a threat to the health and vitality of the vegetation present, and the long term impact of such an introduction on the native vegetation is unknown. It is therefore recommended that the precautionary principle should be adopted for the management of pathogens within all City parks and reserves.






Figure 5 - Pathogen Risk within City of Joondalup Natural Areas.

2.4 High Resolution Multi-Spectral Imagery

As part of the Pathogen Desktop Risk Analysis, the entire extent of the City of Joondalup was acquired with 0.5m high resolution multi-spectral imagery. This involves aerial photography being acquired through a fly over of the area. The imagery includes a near-infrared spectral band that is sensitive to changes in chlorophyll and cell structure of vegetation and can be utilised to assess changes in the vigour and condition of trees and plants over time.

The imagery acquired provides baseline information on the City's vegetation and can be used to monitor the ongoing condition of the City's parks and natural areas to determine areas that may be affected by pathogens by identifying declining health of the vegetation. Variations in vigour are shown as differences in the intensity of red as seen in Figure 7. The imagery can also be a valuable tool in assessing improvement in vegetation vigour for parks and natural areas that have been confirmed as having pathogens present and have had treatment applied.

The imagery also has a number of applications other than for pathogen management. For example the imagery can be utilised to provide baseline information on the extent of the City's vegetation in terms of canopy cover and can be used to monitor the progress of rehabilitation or weed management within parks and natural areas.

It is recommended that the City acquires the high resolution multi-spectral imagery on a biennial basis to enable the ongoing monitoring of the condition of vegetation within the City.



Figure 7- Multi-spectral imagery applied to Granadilla Park.

3.0 Pathogen Management

At present there is no reliable mechanism for the complete eradication of *Phytophthora* species and the control of *Armillaria luteobubalina* is both expensive and labour intensive. Management strategies need to be developed to protect biodiversity values and minimise the spread of these pathogens within the City.

The aim of pathogen management is to:

- Protect biodiversity values within the City with a particular focus on the structure and function of susceptible ecosystems;
- Protect social and economic assets within City parks and natural areas;
- Minimise the spread of pathogens from infected areas to areas that are free from pathogens; and
- Raise community awareness of pathogens and the risk to environmental, social and economic values within the City of Joondalup.

Management strategies should be employed to reduce the spread of pathogens within the City, and initiatives developed to engage the community in order to raise the awareness of pathogen management within the City.

Protecting the biodiversity values of the City's significant bushland areas is a key objective of pathogen management, therefore priority for investigation and the implementation of management recommendations should be given to the City's Major Conservation Areas being:

- Lilburne Park;
- Warwick Open Space;
- Hepburn Conservation Area;
- Shepherd's Bush Park; and
- Craigie Open Space.

The following section provides details of the strategies that should be implemented over the life of the Pathogen Management Plan in order to effectively manage the threat of pathogens within the City of Joondalup.



3.1 Pathogen Interpretation and Mapping

Establishing the extent of Pathogens within the City is essential to minimising the spread of *Phytophthora* species and *Armillaria luteobubalina* within parks and natural areas. Whilst initial desktop investigations have been undertaken to inform the development of this Plan, as detailed in Section 2.3, further investigation is required to ascertain the actual extent of pathogens within the City.

Ground truthing and soil sampling involves site assessments of the park or natural area by a qualified consultant to survey the area for signs of pathogen infestation. Soil and plant samples are also taken from the site which are analysed for the presence of pathogens.

Site investigations to determine the presence of pathogens will enable appropriate responses to effectively manage parks and natural areas where pathogens are confirmed to be present.

As ground truthing and soil sampling can be resource intensive it is recommended that a staged approach is taken to the on ground investigations. It is recommended that a schedule for site investigations is established with parks and natural areas being assessed according to the assigned priority (commencing with Priority 1 reserves) as detailed in Appendix 3.

Once the program for onsite investigations of pathogens has commenced and results of analysis are available it is crucial that this information is communicated to relevant internal and external stakeholders to ensure management actions are applied within the park or natural area.

It is recommended that mapping and a spatial layer be established within the City's Geographic Information System (GIS) to provide information on the areas within the City that are known to have pathogens present.

Management Recommendation 1:

Using Reserve Priority Rankings establish a program for undertaking ground truthing and soil sampling at City parks and natural areas to determine the extent of pathogens within the City of Joondalup.

Management Recommendation 2:

Develop a spatial layer on the City's Geographic Information System (GIS) to provide information to City staff regarding areas of known pathogen infestation.

3.2 Bushland Management

Operational activities of managing bushland reserves have the potential to spread pathogens from one area to another. Pathogens can be spread in small quantities of soil, such as mud attached to tyres, tools or walking shoes.

In order to minimise the risk of spreading pathogens a number of factors should be considered when planning and carrying out bushland management activities. This includes timing of activities such as fire break maintenance, slashing and weeding during dry soil conditions. It is essential that soil movement is minimised and that any materials, tools and vehicles that are bought onto the site are free of pathogens.

Other activities undertaken by the City also risk introducing pathogens to a previously uninfected area, or increase its rate of spread, these include road and drain construction and maintenance. Major land developments, landscaping activities and earthworks which involve the importation or movement of soil and plant material pose significant threats for the introduction and spread of the pathogens throughout the City.

Clear and consistent guidance should be provided to staff and community groups that undertake bushland management activities within vegetated areas of the City in order to minimise the spread of pathogens and prevent new infestations.

Management Recommendation 3:

Develop and implement City of Joondalup Bushland Management Manual to provide staff and community groups with guidance for activities undertaken within parks and natural areas to minimise the spread of pathogens.

3.3 Hygiene Measures

Whether or not pathogens have been identified within a reserve, it is necessary to manage vegetated areas for *Phytophthora* and *Armillaria* in order to prevent its introduction to, or spread within parks and natural areas. Hygiene and quarantine measures are essential to managing pathogens as prevention is much more effective and economical than controlling a pathogen once it is established within a site. The movement of soil and plant parts, such as in plant based mulch from one site to another should be avoided unless it has been confirmed as pathogen free.

Tools, machinery, vehicles and boots should be cleaned of soil after being used within areas of known or suspected infestation. City officers and contractors should also avoid working within infested sites during or immediately after rain or irrigation.

Fire management can also contribute to the introduction and spread of pathogens. Fire prevention activities, such as firebreak construction, can spread infested soil. The Implementation of hygiene procedures when undertaking fire management activities can lead to increased protection of natural areas.

The supply of plant stock, soil and mulch that contains pathogens is one of the most common ways that pathogens such as *Phytophthora* and *Armillaria luteobubalina* establish within an area. It is recommended that the City only purchases plant stock from nurseries accredited under the Nursery Industry Accreditation Scheme Australia (NIASA) and that mulch, soil and gravel supplied for landscaping activities is pathogen free.

Modification of operations that have a high risk of introducing or spreading Phytophthora and Armillaria is also required, particularly when these operations are to occur at sites in or adjacent to areas of high conservation value. The development of guidelines and procedures outlining hygiene measures for City staff, contractors and community groups using City parks and natural areas is essential to controlling the spread of pathogen within the City.

The provisions within the City's Hygiene Guidelines and Management Procedures may also be applicable to the control of other plant diseases within the City including Sting Nematodes.

Management Recommendation 4:

Develop and implement Pathogen Hygiene Guidelines and Management Procedures to provide direction to staff and contractors working within the City's parks and natural areas in order to limit the spread of pathogens within the City of Joondalup.

Management Recommendation 5:

Develop and implement Purchasing Guidelines for the Supply of Landscaping Materials to provide information to City staff, contractors and community groups with information relating to the purchase of plant stock, soil and mulch for City parks and natural areas.

3.4 Access

Many of the City's parks and natural areas have unrestricted access and are often traversed by numerous tracks and paths, increasing the likelihood of pathogens being introduced or spread in the area.

Vehicle access for City parks and natural areas should be restricted, particularly for areas that are known or suspected as having pathogens present. Visitors should be encouraged to avoid using these areas during wet conditions. Visitors should always use tracks and pathways and avoid passing through vegetated areas. Tracks and pathways should be well maintained and have effective drainage. The establishment of new tracks within areas of known infestation should be avoided.

Quarantine measures should be adopted for areas that are known or suspected of being infected by *Phytophthora* or *Armillaria*. Access to these areas by City staff, contractors and the community should be restricted with signage being placed at the entrance of the site explaining the disease status of the park or natural area. Signage should also provide information on how visitors can minimise the spread of the disease within the area.

For high risks areas or areas where Pathogens are known or suspected, cleaning stations should be installed to ensure that footwear, vehicles and tools are free of soil when entering and exiting infected parks or natural areas.

Management Recommendation 6:

Develop a Visitors Guide for City staff, contractors and community groups to provide information regarding access to City parks and natural areas that are known or suspected of having pathogens present.

Management Recommendation 7:

Establish cleaning stations and informational signage at the entrance of City parks and natural areas that are known or suspected of having pathogens present.

3.5 Signage

Communicating the disease status of an area to visitors is an important part of preventing and managing the spread of pathogens. In order to provide a consistent approach to signage within the City it is proposed that the *Western Australian Dieback Signage System* is adopted for use within parks and natural areas. The Signage System was developed by the State Government in partnership with key industry groups as part of a state wide Communication Plan that aims to provide consistent and relevant themes and messages in relation to the management of Phytophthora Dieback within Western Australia.


The Signage System contains a range of signs that can be used depending on the management objectives of the area. The different signs that make up the Signage System are detailed in the *Western Australian Dieback Signage Protocol*, which is included in Appendix 4. The signage utilises a standard set of symbols as illustrated in Figure 8.

Following assessment of a park or natural area by a qualified consultant, the disease status of the area will then be identified as either:

- Pathogen Free
- Pathogen Infested
- Pathogen Unknown



Figure 8- Signage System Disease Status Symbols.



The Signage System also provides information to visitors to assist in minimising the spread of pathogens. Typically signage would be placed at the entrances of the park or natural area with markers being used along paths and tracks.

It is proposed that the term *Pathogen* be used in place of *Dieback* for signage within the City of Joondalup in order to address all four species of *Phytophthora* as well as *Armillaria luteobubalina* in accordance with the objectives of the Pathogen Management Plan.

Management Recommendation 8:

Following site assessment and establishment of the disease status implement the Western Australian Dieback Signage System within City of Joondalup parks and natural areas.

3.6 Communication and Education

Raising the awareness of pathogens within the City of Joondalup is critical to minimising the introduction and spread of disease within vegetated areas of the City. In order to provide ongoing and consistent messages to the community regarding the importance of pathogen management an awareness campaign will be developed and implemented as part of the City's Think Green – Environmental Education Program.

The objectives of the awareness campaign will be to:

- Increase the level of awareness of pathogen management within the community;
- Identify attitudes and behaviours of community members; and
- Develop and implement communication tools that are effective in encouraging measurable changes in behaviour regarding pathogen management within City parks and natural areas.

The campaign will provide information on what pathogens are, how they can be spread and ways of managing the diseases. The program will target the general community, environment groups and schools.

Training of City parks and natural areas staff and members of local environment groups that work within vegetated areas will assist to reduce the risk of introducing and spreading pathogens within the City.

Regular training will ensure that staff and community members are kept up to date with industry developments regarding pathogen management and will include topics such as:

- Communicating outcomes of monitoring programs;
- Hygiene and cleaning requirements;
- Quarantine measures; and
- Treatment options.



Management Recommendation 9:

As part of the City's Think Green Environmental Education Program, develop and implement initiatives that raise the awareness of pathogen management within the community.

Management Recommendation 10:

Ensure City staff and community groups who work within City parks and natural areas undertake relevant training to increase awareness of pathogen control measures.

3.7 Partnerships

Keeping up to date with developments in the area of pathogen management will ensure that the City is implementing best practices approaches to the management of *Phytophthora* and *Armillaria* species. There are a number of private companies, research organisations and working groups within Western Australia that focus on building the capacity of land managers to effectively address pathogens, including the Dieback Information Group and Centre for Phytophthora Science & Management at Murdoch University. The City should actively partner with these organisations to participate in research projects and take up opportunities for sharing information related to best practice approaches to the management of pathogen species.

Management Recommendation 11:

Investigate opportunities to partner with industry groups and research institutions to enable the City to build capacity and gain information relating to best practice approaches to pathogen management.

3.8 Treatment

Once on site investigations have determined that pathogens are present within a park or natural area treatment options should be investigated. Treatment may include removal of infected parts of the specimen, modification of the growing environment and chemical treatments such as Phosphite and growth enhancers.

Phosphite (Phosphonate) is a biodegradable, inexpensive chemical that is systemically transmitted throughout treated plants and has a very low toxicity to animals. Whilst Phosphite does not kill the pathogen, it acts as a booster to native plants enabling them to fight off the pathogen for a significantly longer period of time. Phosphite can be applied ahead of an advancing *Phytophthora* front to form a protective barrier. It can also be applied in an already infested area to protect susceptible plants that have not yet been infected.

Phosphite needs to enter a plant's water transport system in order for it to be effective. This can be done by injecting phosphite directly into the trunk (or roots) of trees, or spraying the leaves of plants.

It is important to note that there is no chemical that

will eradicate *Phytophthora* species and *Armillaria luteobubalina*. However, an integrated approach can successfully control the spread and impact of the diseases. An integrated approach may combine strategic chemical treatment, controlling access, maximising drainage and implementing hygiene and quarantine measures.

As treatment options for *Phytophthora* species and *Armillaria luteobubalina* are both expensive and labour intensive, preventing the introduction of the diseases

into vegetated areas is the main objective of pathogen management.

Management Recommendation 12:

Apply appropriate treatment options for parks and natural areas that have been confirmed to have pathogens present. Investigate and trial newly developed treatment options to determine the best approach to treating pathogens within the City.

4.0 Management Plan Review

The City of Joondalup Pathogen Management Plan will be reviewed and reported on an annual basis to track the progress of the implementation of recommended management actions.

A major review will be undertaken in 2015/16 in order to ensure that the City is managing pathogens in accordance with best practice approaches.



5.0 References

Barber, P - Arbor Carbon (2012) Desktop Risk Analysis of Phytophthora and Armillaria to Reserves Within the City of Joondalup, Perth Western Australia

Dieback Working Group (2000) Managing Dieback Guidelines for Local Government, Perth Western Australia

Dieback Working Group (2008) Standart Dieback Signage Protocol for use, Perth WA.

Dieback Working Group (2008) Managing Phytophthora Dieback in Bushland, Perth Western Australia

Forest Science Centre (2003) Ian W. Smith & David I. Smith, Armillaria Root Rot: A Disease of Native and Introduced Trees, Victoria Australia

Smith-White J.L and Summerell B.A (2003) Armillaria Root Rot- Under the Microscope, Microbiology Australia


Project Dieback (2008) Standard Dieback Signage Protocol, Western Australia

Shearer, B.L., Crane, C.E. and Cochrane, A. (2004) *Quantification of the susceptibility of the native flora of the southwest botanical province, Western Australia, to Phytophthora cinnamomi*, Australian Journal of Botany. 52: 435-443

Shearer, B. L and Tippett J. T (1988) *Distribution and Impact of Armillaria luteobubalina in the Eucalyptus marginata Forest of South-Western Australia*, Australian Journal of Botany 36(4) 433 – 445

University of Western Australia (2004) Soil Health- Development and Prevention of Plant Disease www.soilhealth.see.uwa.edu.au

Western Australian Local Government Association (2004) Local Government Biodiversity Planning Guidelines, Western Australia



6.0 Appendices

- Appendix 1 Sites included in Pathogen Risk Assessment Study
- Appendix 2 Pathogen Desktop Risk Assessment Criteria
- Appendix 3 Priority List of Parks and Natural Areas for Pathogen Mapping and Monitoring
- Appendix 4 Western Australian Dieback Signage Protocol

Appendix 1

List of Sites within Study Area

Site Name	Address	Suburb
Alidade Park	Alidade Way	Beldon
Beldon Park	Gradient Way	Beldon
Cumberland Park	Cumberland Way	Beldon
Ensign Park	Ensign Way	Beldon
Gradient Park	Gradient Way	Beldon
Haddington Park	Haddington Street	Beldon
Monument Park	Monument Drive	Beldon
Sandalford Park	Sandalford Drive	Beldon
Beachside Park	Beachside Drive	Burns Beach
Bramston Park	Bramston Vista	Burns Beach
Burns Beach Park	Ocean Parade	Burns Beach
Burns Park	Ocean Parade	Burns Beach
Coastal Foreshore Reserve Burns Beach		Burns Beach
Huxley Park	Huxleys Trail	Burns Beach
Southport Park	Southport Loop	Burns Beach
Third Park	Third Avenue	Burns Beach
Windmill Park	Windmill Circle	Burns Beach
Baltusrol Park	Baltusrol Rise	Connolly
Bonnie Doon Park	Bonnie Doon Gardens	Connolly
Carnaby Reserve	Medinah Mews	Connolly
Cinque Ports Park	Cinque Ports Place	Connolly
Fairway Park	Fairway Circle	Connolly
Huntingdale Park	Huntingdale Crescent	Connolly Connolly
Pine Valley Park St Michael's Park	Pine Valley Pass St Michael'S Avenue	Connolly
		-
Adelaide Park	Adelaide Circle	Craigie
Albion Park	Camberwarra Drive	Craigie
Barwon Park	Barwon Road	Craigie
Camberwarra Park	Camberwarra Drive	Craigie
Cawarra Park	Cawarra Crescent	Craigie
Chadstone Park	Chadstone Road	Craigie
Craigie Open Space	Whitfords Avenue	Craigie
Craigie Park	Barradine Way	Craigie
Madana Park	Madana Place	Craigie
Mandalay Park	Mandalay Place	Craigie
Otago Park	Camberwarra Drive	Craigie
Otway Park	Camberwarra Drive	Craigie
Warrandyte Park	Warrandyte Drive	Craigie
Whitfords East Park	Rosette Close	Craigie
Caledonia Park	Caledonia Avenue	Currambine
Carlton Park	Carlton Turn	Currambine
Christchurch Park	Christchurch Terrace	Currambine
Clermont Park	Caledonia Avenue	Currambine
De Crillon Park	De Crillon Way	Currambine
Doncaster Park	Doncaster Square	Currambine

Site Name	Address	Suburb
Fairmont Park	Fairmont Place	Currambine
Negresco Park	Negresco Turn	Currambine
Riversdale Park	Riversdale Gardens	Currambine
Santa Ana Park	Aberfoyle Heights	Currambine
Alder Park	Alder Way	Duncraig
Alfreton Park	Alfreton Way	Duncraig
Bracken Park	Bracken Court	Duncraig
Buckthorn Park	Buckthorn Way	Duncraig
Castlefern Park	Castlefern Way	Duncraig
Coolibah Park	Coolibah Place	Duncraig
Galston Park	Galston Place	Duncraig
Geddes Park	Geddes Close	Duncraig
Glenbar Park	Glenbar Road	Duncraig
Glengarry Park	Merrick Way	Duncraig
Granadilla Park	Langholm Place	Duncraig
Greenlaw Park	Greenlaw Street	Duncraig
Hilton Park	Hilton Place	Duncraig
Juniper Park	Juniper Way	Duncraig
Kelvin Park	Campion Street	Duncraig
Lanark Park	Lanark Mews	Duncraig
Lilburne Park	Lilburne Road	Duncraig
Macaulay Park	Macaulay Avenue	Duncraig
Marri Park	Marri Road	Duncraig
Melene Park	Melene Road	Duncraig
Noel Gannon Park	Lennoxtown Road	Duncraig
Oakapple Park	Oakapple Drive	Duncraig
Pentland Park	Pentland Avenue	Duncraig
Percy Doyle Reserve	Warwick Road	Duncraig
Portree Park	Portree Way	Duncraig
Poynter Park	Poynter Drive	Duncraig
Salata Park	Salata Place	Duncraig
Sycamore Park	Scaddan Street	Duncraig
Telopia Park	Telopia Drive	Duncraig
Trigonometric Park	Truslove Way	Duncraig
Wanbrow Park	Wanbrow Way	Duncraig
Wandina Park	Wandina Place	Duncraig
Beaumont Park	Beaumont Way	Edgewater
Edgewater Park	Edgewater Drive	Edgewater
Emerald Park	Emerald Way	Edgewater
Garrong Park	Garrong Close	Edgewater
-	Hilltop Place	
Hilltop Park Lakevalley Park	Lakevalley Drive	Edgewater Edgewater
Site Name	Address	Suburb
Lakeview Park	Lakeview Drive	
		Edgewater
Lookout Park	Quarry Ramble	Edgewater
Okely Park	Joondalup Drive	Edgewater
Picnic Cove Park	Edgewater Drive	Edgewater
Quarry Park	Joondalup Drive	Edgewater
Quarry Ramble Park	Quary Ramble	Edgewater
Ridge Park	Ridge Close	Edgewater

Site Name	Address	Suburb
St Clair Park	Joondalup Drive	Edgewater
Stillwater Park	Reflection Close	Edgewater
Tuart Park	Tuart Trail	Edgewater
Wedgewood Park	Wedgewood Drive	Edgewater
Annato Park	Annato Street	Greenwood
Birch Park	Birch Place	Greenwood
Blackall Park	Blackall Drive	Greenwood
Blackthorn Park	Blackthorn Road	Greenwood
Calectasia Park	Calectasia Street	Greenwood
Cockman Park	Cockman Road	Greenwood
Filbert Park	Filbert Street	Greenwood
George Sears Park	Wanneroo Road	Greenwood
Gerda Park	Gerda Court	Greenwood
Hartley Park	Hartley Court	Greenwood
Kanangra Park	Kanangra Crescent	Greenwood
Karuah Park	Karuah Way	Greenwood
Kilrenny Park	Gilmerton Way	Greenwood
Kurrajong Park	Kurrajong Place	Greenwood
Leaside Park	Leaside Way	Greenwood
Mamo Park	Mamo Place	Greenwood
Oleaster Park	Oleaster Way	Greenwood
Parin Park	Canham Way	Greenwood
Paveta Park	Paveta Court	Greenwood
Penistone Park	Penistone Street	Greenwood
Rodgers Park	Rodgers Street	Greenwood
Sheoak Park	Tuart Road	Greenwood
Sherington Park	Sherington Road	Greenwood
Strathaven Park	Strathaven Crescent	Greenwood
Warrigal Park	Madrona Crescent	Greenwood
Abrolhos Park	Abrolhos Drive	Heathridge
Admiral Park	Admiral Grove	Heathridge
Balanus Park	Balanus Way	Heathridge
Conidae Park	Conidae Drive	Heathridge
Faversham Park	Faversham Way	Heathridge
Heathridge Park	Sail Terrace	Heathridge
Larkspur Park	Larkspur Place	Heathridge
Littorina Park	Littorina Avenue	Heathridge
Lysander Park	Lysander Drive	Heathridge
Poseidon Park	Poseidon Road	Heathridge
Prince Regent Park	Prince Regent Drive	Heathridge
Banks Park	Banks Avenue	Hillarys
Broadbeach Park	Broadbeach Boulevard	Hillarys
Conica Park	Conica Crescent	Hillarys
Fenton Park	Fenton Way	Hillarys
Flinders Park	Centennial Gardens	Hillarys
Gleddon Park	Gleddon Way	Hillarys
Harbour View Park	Azzuro Crescent	Hillarys
Harcourt Park	Harcourt Drive	Hillarys
Hillarys Park	Absolon Way	Hillarys
James Cook Park	Banks Avenue	Hillarys
		- · , -

Site Name	Address	Suburb
Leeward Park	Leeward Chase	Hillarys
Lymburner Park	Lymburner Drive	Hillarys
Maquire Park	Maquire Road	Hillarys
Marbella Park	Mallorca Avenue	Hillarys
Mawson Park	Mawson Crescent	Hillarys
Nash Park	Nash Street	Hillarys
Nimrod Park	Nimrod Place	Hillarys
Ninnis Park	Ninnis Place South	Hillarys
Oahu Park	Oahu Gardens	Hillarys
Orient Park	Orient Circuit	Hillarys
Parkinson Park	Parkinson Place	Hillarys
Pinnaroo Point	Whitfords Avenue	Hillarys
Scott Park	Webb Place	Hillarys
Whitford Nodes Central	Whitfords Avenue	Hillarys
Whitford Nodes South	Whiterds Avenue	Hillarys
Whitford Nodes South	White Avenue	Hillarys
Bethany Park	Bethany Gardens	Iluka
Coastal Foreshore Reserve Iluka	Burns Beach Road	Iluka
Discovery Park	Discovery Circuit	Iluka
Iluka District Open Space	Miami Beach Promenade	Iluka
Iluka Foreshore Reserve	Ocean Reef Road	Iluka
Kuta Park	Kuta Circuit	Iluka
Naturaliste Park	Naturaliste Boulevard	lluka
Sir James McCusker Park	Silver Sands Drive	lluka
Albright Park	Cornell Parade	Joondalup
Aldgate Park	Aldgate Street	Joondalup
Blue Lake Park	Blue Mountain Drive	Joondalup
Candlewood Park	Candlewood Boulevard	Joondalup
Central Park	Grand Boulevard	Joondalup
Charing Cross Park	Charing Cross Road	Joondalup
Elcar Park		Joondalup
Embankment Park	The Embankment	Joondalup
Greenshank Park	Cockatoo Ridge	Joondalup
Katrine Park	Katrine Crescent	Joondalup
Lady Evelyn Park	Lady Evelyn Retreat	Joondalup
Lakeside Park	Waterbird Turn	Joondalup
Manapouri Park	Manapouri Meander	Joondalup
Nanika Park	Nanika Crescent	Joondalup
Neil Hawkins Park	Boas Avenue	Joondalup
Piccadilly Park	Piccadilly Circle	Joondalup
Portwood Park	Cornell Parade	Joondalup
Queensbury Park	Queensbury Road	Joondalup
Regents Park	Regents Park Road	Joondalup
Sittella Park	Sittella Turn	Joondalup
Stilt Park	Cockatoo Ridge	Joondalup
Thornbill Park	Thornbill Meander	Joondalup
Walsh Park	Joondalup Drive	Joondalup
Water Tower Park	Moondarra Way	Joondalup
Wesley Park	Cornell Parade	Joondalup
Windermere Park	Candlewood Boulevard	Joondalup

Site Name	Address	Suburb
Woodlea Park	Woodlea Crescent	Joondalup
Aristride Park	Aristride Avenue	Kallaroo
Belrose Park	Belrose Entrance	Kallaroo
Bridgewater Park	Bridgewater Drive	Kallaroo
Castlecrag Park	Castlecrag Drive	Kallaroo
Culwalla Park	Culwalla Close	Kallaroo
Dampier Park	Dampier Avenue	Kallaroo
Delaware Park	Delaware Place	Kallaroo
Glenbank Park	Awhina Place	Kallaroo
Kiernan Park	Kiernan Place	Kallaroo
Kilarney Park	Kilarney Heights	Kallaroo
Maritana Park	Maritana Road	Kallaroo
Merrifield Park	Northshore Drive	Kallaroo
Montague Park	Montague Way	Kallaroo
Stanford Park	Stanford Road	Kallaroo
Whitford Nodes North	Northshore Drive	Kallaroo
Whitfords West Park	Marmion Avenue	Kallaroo
Barridale Park	Barridale Drive	Kingsley
Bindaree Park	Bindaree Terrace	Kingsley
Cadogan Park	Cadogan Street	Kingsley
Calthorpe Park	Calthorpe Place	Kingsley
Chelsea Park	Chelsea Court	Kingsley
Dollis Park	Dollis Way	Kingsley
Forest Hill Park	Kingsley Drive	Kingsley
Greenwich Park	Greenwich Court	Kingsley
Illawong Park	Illawong Way	Kingsley
Kingfisher Park	Kingfisher Way	Kingsley
Kingsley Park	Kingsley Drive	Kingsley
Legana Park	Legana Avenue	Kingsley
Lehmann Park	Lehmann Court	Kingsley
Moolanda Park	Moolanda Boulevard	Kingsley
Newham Park	Newham Way	Kingsley
Plover Park	Plover Way	Kingsley
Robertson Road Cycleway	Barridale Drive	Kingsley
Shepherds Bush Park	Shepherds Bush Drive	Kingsley
Spoonbill Park	Spoonbill Grove	Kingsley
Talbot Park	Talbot Drive	Kingsley
Wallangarra Park	Wallangarra Court	Kingsley
Callander Park	Callander Avenue	Kinross
Cranston Park	Cranston Loop	Kinross
Earlsferry Park	Earlsferry Green	Kinross
Falkland Park	Falkland Way	Kinross
MacNaughton Park	Macnaughton Cresecent	Kinross
Menteith Park	Selkirk Drive	Kinross
Roxburgh Park	Roxburgh Circle	Kinross
Rutherglen Park	Rutherglen Circle	Kinross
Stonehaven Park	Stonehaven Parade	Kinross
Stow Park	Stow Lane	Kinross
Thornton Park	Thornton Retreat	Kinross
Banksia Park	Banksia Dale	Marmion
Dai Insia Fai K		

Address	Suburb
Braden Way	Marmion
Cliff Street	Marmion
Troy Avenue	Marmion
Cliverton Court	Marmion
West Coast Drive	Marmion
Finney Crescent	Marmion
Keppell Road	Marmion
Ozone Road	Marmion
Mckirdy Way	Marmion
Anemone Way	Mullaloo
-	Mullaloo
Charonia Road	Mullaloo
Gunida Way	Mullaloo
Kallaroo Place	Mullaloo
Korella Street	Mullaloo
Periwinkle Road	Mullaloo
Oceanside Promenade	Mullaloo
Triton Place	Mullaloo
	Suburb
	Mullaloo
	Mullaloo
	Ocean Reef
	Padbury
-	Padbury
	-
Newcombe Way	Padubry
Newcombe Way Santiago	Padubry Parkway
	Braden WayCliff StreetTroy AvenueCliverton CourtWest Coast DriveFinney CrescentKeppell RoadOzone RoadMckirdy WayAnemone WayBalga WayCharonia RoadGunida WayKallaroo PlaceKorella StreetPeriwinkle RoadOceanside Promenade

Site Name	Address	Suburb
Clare Park	Clare Street	Sorrento
Geneff Park	Padbury Circle	Sorrento
Hakea Park	Hakea Place	Sorrento
Harman Park	Harman Road	Sorrento
Lacepede Park	Lacepede Drive	Sorrento
Porteous Park	Porteous Road	Sorrento
Robin Park	Robin Avenue	Sorrento
Seacrest Park	Seacrest Drive	Sorrento
Sorrento Foreshore	West Coast Drive	Sorrento
Tom Walker Park	Sandpiper Street	Sorrento
Aberdare Park	Aberdare Way	Warwick
Carr Park	Carr Crescent	Warwick
Chelsford Park	Chelsford Road	Warwick
Churton Park	Churton Crescent	Warwick
Circle Park	The Circle	Warwick
Ellersdale Park	Ellersdale Avenue	Warwick
Glenmere Park	Glenmere Road	Warwick
Hawker Park	Hawker Avenue	Warwick
Hillwood Park (North)	Hillwood Avenue	Warwick
Hillwood Park (South)	Hillwood Avenue	Warwick
Springvale Park	Springvale Drive	Warwick
Warwick Open Space	Beach Road	Warwick
Chichester Park	Trappers Drive	Woodvale
Cornish Park	Cornish Avenue	Woodvale
Delonix Park	Delonix Circle	Woodvale
Gascoyne Park	Gascoyne Avenue	Woodvale
McCubbin Park	Mccubbin Boulevard	Woodvale
Parkside Park	Parkside Ramble	Woodvale
Plumdale Park	Plumdale Way	Woodvale
Timbercrest Park	Trailwood Drive	Woodvale
Timberlane Park	Althaea Way	Woodvale
Trailwood Park	Trailwood Drive	Woodvale
Trappers Park	Trappers Drive	Woodvale
Waterview Park	Waterview Drive	Woodvale

Appendix 2

Pathogen Desktop Risk Assessment Criteria

Parks Assessment - Part 1: Disease Risk

Variable	Calculation
Disease confirmed in or within a connected park	50
Disease suspected in or within a connected park	35
Disease not confirmed or suspected in or within a connected park	0
Connected to a Natural Area	25
Not connected to a Natural Area	0
Irrigated	10
Not irrigated	0

Table 1- The variables used and their associated values inserted into the calculation to determine the total risk value and ranking of Parks in order of risk of disease from the threatening pathogens.

An example of the calculation used for a typical park such Aberdare Park, which had no confirmation of disease or suspected disease in or within a connected park, is not connected to a Natural Area, and is irrigated, would be as follows:

0 + 0 + 10 = 10

Parks ranged in total values from 71 down to 0. The Disease Risks for the parks was determined using the above criteria and categorised as either:

- Risk category 1 (High) greater than 10
- Risk category 2 (Medium) 1 to 10
- Risk category 3 (Low) 0

Parks Assessment - Part 2: Overall Risk and Prioritisation for Pathogen Monitoring and Mapping

Variable	Calculation
Disease Risk Category High	15
Disease Risk Category Medium	10
Disease Risk Category Low	5
Heddle Vegetation Complex	25
Karrakatta complex central and south	4
Herdsman complex	3
Cottesloe complex central and south	2
Quindalup complex	1

Table 2- The variables used and their associated values inserted into the calculation to determine ranking of parks for overall risk and prioritisation of Phase 2 activities.

An example of the calculation used for a typical park such as Beachside Park, which had a Disease Risk ranking of medium, and is comprised of vegetation within the Quindalup complex, would be as follows:

10 + 1 = 11

Parks ranged in total values from 19 down to 5. The overall risk and priority for Phase 2 assessment (Pathogen Mapping and Monitoring) was determined using the above criteria and categorised as either:

- Priority 1 (High) greater than 10
- Priority 2 (Medium) 10
- Priority 3 (Low) 5 to 9

Natural Areas Assessment - Part 1: Disease Risk

Variable	Calculation
Disease confirmed in or within a Natural Area	50
Disease suspected in or within a Natural Area	35
Disease not confirmed or suspected in or within a connected Natural Area/Park	0
Connected to a Bush Forever Park	1
Not connected to a Bush Forever Park	0
Connected to an irrigated park	1
Not connected to an irrigated park	0

Table 3- The variables used and their associated values inserted into the calculation to determine the total risk value and ranking of Natural Areas in order of risk of disease from the threatening pathogens.

An example of the calculation used for a typical Natural Area such Cadogan Park, which had no confirmation of disease but was suspected of having disease, is not connected to a Bush Forever site, and is not irrigated, would be as follows:

35 + 0 + 0 = 35

Natural Areas ranged in total values from 52 down to 0. The Disease Risks for the natural areas was determined using the above criteria and categorised as either:

- Risk category 1 (High) > 34
- Risk category 2 (Medium) 1 to 34
- Risk category 3 (Low) 0

Natural Areas Assessment - Part 2: Overall Risk and Prioritisation for Pathogen Monitoring and Mapping

Variable	Calculation
Disease Risk Category High	150
Disease Risk Category Medium	100
Disease Risk Category Low	50
Heddle Vegetation Complex	
Karrakatta complex central and south	4
Herdsman complex	3
Cottesloe complex central and south	2
Quindalup complex	1
NAIA Priority Rank	100 minus rank value

Table 4- The variables used and their associated values inserted into the calculation to determine ranking of natural areas for overall risk and prioritisation of Phase 2 activities

An example of the calculation used for a typical Natural Area such as Craigie Open Space, which had a Disease Risk ranking of high, and is comprised of vegetation within the Karrakatta complex central and south, and a NAIA Priority ranking of 4 would be as follows:

150 + 4 + 96 = 250

Natural Areas ranged in total values from 250 down to 50. The overall risk and priority for Phase 2 assessment (Pathogen Mapping and Monitoring) was determined using the above criteria and categorised as either:

- Priority 1 (High) -> 149
- Priority 2 (Medium) 100 to 149
- Priority 3 (Low) 50 to 99

Appendix 3

Overall Risk / Priority Listings for Parks and Natural Areas

Priority 1 - Parks and Natural Areas

Name	Address	Suburb
Aberdare Park	Aberdare Way	Warwick
Baltusrol Park	Baltusrol Rise	Connolly
Beachside Park	Beachside Drive	Burns Beach
Beaumaris Park	Beaumaris Boulevard	Ocean Reef
Blackboy Park	Balga Way	Mullaloo
Blue Lake Park	Blue Mountain Drive	Joondalup
Bonnie Doon Park	Bonnie Doon Gardens	Connolly
Brazier Park	Brazier Rise	Padbury
Burns Beach Park	Ocean Parade	Burns Beach
Burns Beach Park	Ocean Parade	Burns Beach
Cadogan Park	Cadogan Street	Kingsley
Central Park	Lakeside Drive	Joondalup
Chichester Park	Trappers Drive	Woodvale
Clermont Park	Caledonia Avenue	Currambine
Craigie Open Space	Whitfords Avenue	Craigie
Cranston Park	Cranston Loop	Kinross
Emerald Park	Emerald Way	Edgewater
Fairway Park	Fairway Circle	Connolly
Fernwood Park	Fernwood Square	Padbury
Garrong Park	Garrong Close	Edgewater
Gibson Park	Gibson Avenue	Padbury
Granadilla Park	Granadilla Street	Duncraig
Greenshank Park	Cockatoo Ridge	Joondalup
Hawker Park	Hawker Avenue	Warwick
Hepburn Conservation Area	Hepburn Avenue	Padbury
Hillarys Park	Absolon Way	Hillarys
Juniper Park	Juniper Way	Duncraig
Lakeside Park	Waterbird Turn	Joondalup
Lakevalley Park	Lakevalley Drive	Edgewater
Legana Park	Legana Avenue	Kingsley
Lexcen Park	Venturi Drive	Ocean Reef
Lilburne Park	Lilburne Road	Duncraig
Macnaughton Park	Macnaughton Cresecent	Kinross
Maritana Park	Maritana Road	Kallaroo
Merrifield Park	Northshore Drive	Kallaroo
Nanika Park	Nanika Crescent	Joondalup
Naturaliste Park	Naturaliste Boulevard	Iluka
Negresco Park	Negresco Turn	Currambine
Neil Hawkins Park	Boas Avenue	Joondalup
Plumdale Park	Plumdale Way	Woodvale
Quarry Park	Regatta Drive	Edgewater
Robertson Road Cycleway	Barridale Drive	Kingsley
Robin Park	Robin Avenue	Sorrento

Name	Address	Suburb	
Sandalford Park	Sandalford Drive	Beldon	
Shepherds Bush Park	Shepherds Bush Drive	Kingsley	
Sir James Mccusker Park	Silver Sands Drive	Iluka	
St Clair Park	Joondalup Drive	Edgewater	
St Michael's Park	St Michael's Avenue	Connolly	
Third Park	Third Avenue	Burns Beach	
Timberlane Park	Althaea Way	Woodvale	
Tom Simpson Park	Oceanside Promenade	Mullaloo	
Trappers Park	Trappers Drive	Woodvale	
Warrandyte Park	Warrandyte Drive	Craigie	
Warwick Open Space	Beach Road	Warwick	
Waterview Park	Waterview Drive	Woodvale	
Whitford Nodes South	Whitfords Avenue	Kallaroo	

Priority 2 - Parks and Natural Areas

Name	Street	Suburb	
Abrolhos Park	Abrolhos Drive	Heathridge	
Adelaide Park	Adelaide Circle	Craigie	-
Admiral Park	Admiral Grove	Heathridge	
Albacore Park	Albacore Drive	Sorrento	
Albright Park	Cornell Parade	Joondalup	
Aldgate Park	Aldgate Street	Joondalup	
Alfreton Park	Alfreton Way	Duncraig	
Annato Park	Annato Street	Greenwood	
Aristride Park	Aristride Avenue	Kallaroo	
Balanus Park	Balanus Way	Heathridge	
Banks Park	Banks Avenue	Hillarys	
Barridale Park	Barridale Drive	Kingsley	
Barwon Park	Barwon Road	Craigie	
Barwon Park	Barwon Road	Craigie	
Beaumont Park	Beaumont Way	Edgewater	
Beenyup Park	Timerlane Drive	Woodvale	
Beldon Park	Gradient Way	Beldon	
Belrose Park	Belrose Entrance	Kallaroo	
Bethany Park	Bethany Gardens	Iluka	
Bindaree Park	Bindaree Terrace	Kingsley	
Birdland Park	Birdland Court	Edgewater	
Blackall Park	Cedarwood Circle	Greenwood	
Blackall Park	Blackall Drive	Greenwood	
Blackthorn Park	Blackthorn Road	Greenwood	
Blue Lake Park	Blue Mountain Drive	Joondalup	
Braden Park	Braden Way	Marmion	
Bridgewater Park	Bridgewater Drive	Kallaroo	
Brisbane Park	Brisbane Drive	Padbury	
Broadbeach Park	Broadbeach Boulevard	Hillarys	
Buckthorn Park	Buckthorn Way	Duncraig	
Cadogan Park	Cadogan Street	Kingsley	
Caledonia Park	Caledonia Avenue	Currambine	
Callander Park	Callander Avenue	Kinross	_

Name	Address	Suburb
Callion Park	Pinnaroo Drive	Padbury
Calthorpe Park	Calthorpe Place	Kingsley
Camberwarra Park	Camberwarra Drive	Craigie
Carina Park	Carina Loop	Ocean Reef
Carlton Park	Carlton Turn	Currambine
Castlecrag Park	Castlecrag Drive	Kallaroo
Castlefern Park	Castlefern Way	Duncraig
Cawarra Park	Cawarra Crescent	Craigie
Chadstone Park	Chadstone Road	Craigie
Charing Cross Park	Aldwych Way	Joondalup
Charonia Park	Charonia Road	Mullaloo
Chelsea Park	Chelsea Court	Kingsley
Chelsford Park	Chelsford Road	Warwick
Christchurch Park	Christchurch Terrace	Currambine
Churton Park	Churton Crescent	Warwick
Cinque Ports Park	Cinque Ports Place	Connolly
Circle Park	The Circle	Warwick
Coastal Foreshore Reserve Iluka	Burns Beach Road	lluka
Cockman Park	Cockman Road	Greenwood
Conica Park	Conica Crescent	Hillarys
Conidae Park	Conidae Drive	Heathridge
Coolibah Park	Coolibah Place	Duncraig
Cunningham Park	Cunningham Place	Padbury
Dampier Park	Dampier Avenue	Kallaroo
De Crillon Park	De Crillon Way	Currambine
Delaware Park	Delaware Place	Kallaroo
Delonix Park	Delonix Circle	Woodvale
Discovery Park	Discovery Circuit	lluka
Dollis Park	Dollis Way	Kingsley
Doncaster Park	Doncaster Square	Currambine
Earlsferry Park	Earlsferry Green	Kinross
Edgewater Park	Edgewater Drive	Edgewater
Elcar Park	Elcar Lane	Joondalup
Ellersdale Park	Ellersdale Avenue	Warwick
Embankment Park	The Embankment	Joondalup
Ensign Park	Ensign Way	Beldon
Fairmont Park	Fairmont Place	Currambine
Falkland Park	Falkland Way	Kinross
Faversham Park	Faversham Way	Heathridge
Fenton Park	Fenton Way	Hillarys
Finney Park	Finney Crescent	Marmion
Flinders Park	Centennial Gardens	Hillarys
Forest Hill Park	Kingsley Drive	Kingsley
Forrest Park	Forrest Road	Padbury
Galston Park	Galston Place	Duncraig
Gascoyne Park	Gascoyne Avenue	Woodvale
Geddes Park	Geddes Close	Duncraig
Geneff Park	Padbury Circle	Sorrento
	···· , · ···	
George Sears Park	Wanneroo Road	Greenwood

Name	Address	Suburb
Glenbank Park	Awhina Place	Kallaroo
Glenbar Park	Glenbar Road	Duncraig
Glengarry Park	Merrick Way	Duncraig
Gradient Park	Gradient Way	Beldon
Greenlaw Park	Greenlaw Street	Duncraig
Haddington Park	Haddington Street	Beldon
Harbour View Park	Azzuro Crescent	Hillarys
Heathridge Park	Sail Terrace	Heathridge
Herreshoff Park	Herreshoff Ramble	Ocean Reef
Hilltop Park	Hilltop Place	Edgewater
Hilton Park	Hilton Place	Duncraig
Huntingdale Park	Huntingdale Crescent	Connolly
Illawong Park	Illawong Way	Kingsley
Iluka District Open Space	Miami Beach Promenade	lluka
Iluka Foreshore Reserve	Ocean Reef Road	lluka
James Cook Park	Banks Avenue	Hillarys
Kallaroo Park	Kallaroo Place	Mullaloo
Kanangra Park	Kanangra Crescent	Greenwood
Katrine Park	Katrine Crescent	Joondalup
Keppell Park	Keppell Road	Marmion
Kiernan Park	Kiernan Place	Kallaroo
Kilarney Park	Kilarney Heights	Kallaroo
Killin Park	Sycamore Drive	Duncraig
Kimberley Park	Kimberley Road	Hillarys
Kingsley Park	Kingsley Drive	Kingsley
Korella Park	Korella Street	Mullaloo
Lacepede Park	Lacepede Drive	Sorrento
Lacepede Park	Lacepede Drive	Sorrento
Lady Evelyn Park	Lady Evelyn Retreat	Joondalup
Larkspur Park	Larkspur Place	Heathridge
Leeward Park	Leeward Chase	Hillarys
Lehmann Park	Lehmann Court	Kingsley
Leichhardt Park No1	Leichhardt Avenue	Padbury
Littorina Park	Littorina Avenue	Heathridge
Lymburner Park	Lymburner Drive	Hillarys
Lysander Park	Lysander Drive	Heathridge
Macaulay Park	Macaulay Avenue	Duncraig
Macdonald Park	Macdonald Avenue	Padbury
Mamo Park	Mamo Place	Greenwood
Mamo Park	Cedarwood Circle	Greenwood
Manapouri Park	Manapouri Meander	Joondalup
Maquire Park	Maquire Road	Hillarys
Marbella Park	Mallorca Avenue	Hillarys
Marri Park	Marri Road	Duncraig
Mawson Park	Mawson Crescent	Hillarys
Mccubbin Park	Mccubbin Boulevard	Woodvale
Mckirdy Park	Mckirdy Way	Marmion
Malaza Davis		
Melene Park	Melene Road	Duncraig
Melene Park Menteith Park	Melene Road Selkirk Drive	Duncraig Kinross

Name	Address	Suburb
Montague Park	Montague Way	Kallaroo
Moolanda Park	Moolanda Boulevard	Kingsley
Nash Park	Nash Street	Hillarys
Newcombe Park	Newcombe Way	Padubry
Newham Park	Newham Way	Kingsley
Ninnis Park	Ninnis Place South	Hillarys
Noel Gannon Park	Lennoxtown Road	Duncraig
Oahu Park	Oahu Gardens	Hillarys
Ocean Reef Park	Marina Boulevard	Ocean Reef
Oleaster Park	Oleaster Way	Greenwood
Orient Park	Orient Circuit	Hillarys
Otago Park	Camberwarra Drive	Craigie
Oxley Park	Oxley Avenue	Padbury
Parin Park	Canham Way	Greenwood
Parkinson Park	Parkinson Place	Hillarys
Parkside Park	Parkside Ramble	Woodvale
Penistone Park	Penistone Street	Greenwood
Percy Doyle Reserve	Warwick Road	Duncraig
Periwinkle Park	Periwinkle Road	Mullaloo
Piccadilly Park	Piccadilly Circle	Joondalup
Picnic Cove Park	Edgewater Drive	Edgewater
		Connolly
Pine Valley Park	Pine Valley Pass	
Pinnaroo Point	Whitfords Avenue	Hillarys
Porteous Park	Porteous Road	Sorrento
Portree Park	Portree Way	Duncraig
Portwood Park	Lakeside Drive	Joondalup
Poseidon Park	Poseidon Road	Heathridge
Poynter Park	Poynter Drive	Duncraig
Prince Regent Park	Prince Regent Drive	Heathridge
Quarry Ramble Park	Quary Ramble	Edgewater
Queensbury Park	Queensbury Road	Joondalup
Regents Park	Regents Park Road	Joondalup
Rev John Smithies Park	Lakeway Drive	Kingsley
Riversdale Park	Riversdale Gardens	Currambine
Rodgers Park	Rodgers Street	Greenwood
Roxburgh Park	Roxburgh Circle	Kinross
Rutherglen Park	Rutherglen Circle	Kinross
Salata Park	Salata Place	Duncraig
Santiago Park	Santiago	Parkway
Scott Park	Webb Place	Hillarys
Seacrest Park	Seacrest Drive	Sorrento
Sheoak Park	Tuart Road	Greenwood
Sherington Park	Sherington Road	Greenwood
Simpson Park	Forrest Road	Padbury
Sittella Park	Sittella Turn	Joondalup
Sorrento Foreshore	West Coast Drive	Sorrento
Southern Cross Park	Souther Cross Circle	Ocean Reef
Springvale Park	Springvale Drive	Warwick
Stanford Park	Stanford Road	Kallaroo
Stilt Park	Cockatoo Ridge	Joondalup

Name	Address	Suburb	
Stonehaven Park	Stonehaven Parade	Kinross	
Stow Park	Stow Lane	Kinross	
Sycamore Park	Scaddan Street	Duncraig	
Tarolinta Park	Tarolinta Gardens	Ocean Reef	
Telopia Park	Telopia Drive	Duncraig	
Third Park	Third Avenue	Burns Beach	
Thornbill Park	Thornbill Meander	Joondalup	
Thornton Park	Thornton Retreat	Kinross	
Timbercrest Park	Trailwood Drive	Woodvale	
Tom Walker Park	Sandpiper Street	Sorrento	
Trig Point Park	Shoran Court	Ocean Reef	
Trigonometric Park	Truslove Way	Duncraig	
Triton Park	Triton Place	Mullaloo	
Wallangarra Park	Wallangarra Court	Kingsley	
Walsh Park	Joondalup Drive	Joondalup	
Wanbrow Park	Wanbrow Way	Duncraig	
Warrigal Park	Madrona Crescent	Greenwood	
Water Tower Park	Moondarra Way	Joondalup	
Wedgewood Park	Wedgewood Drive	Edgewater	
Wesley Park	Cornell Parade	Joondalup	
Whistler Park	Whistler Close	Edgewater	
Whitfords East Park	Rosette Close	Craigie	
Whitfords West Park	Marmion Avenue	Kallaroo	
Windermere Park	Candlewood Boulevard	Joondalup	
Wolinski Park	Key West Drive	Mullaloo	
Woodlea Park	Woodlea Crescent	Joondalup	

Priority 3- Parks and Natural Areas

Name	Street	Suburb	
Albion Park	Camberwarra Drive	Craigie	
Alder Park	Alder Way	Duncraig	
Alidade Park	Alidade Way	Beldon	
Anemone Park	Anemone Way	Mullaloo	
Bage Park	Bage Court	Hillarys	
Banksia Park	Banksia Dale	Marmion	
Bannister Park	Bannister Road	Padbury	
Barclay Park	Barclay Avenue	Padbury	
Batavia Park	Batavia Place	Kallaroo	
Bigola Park	Bigola Court	Kallaroo	
Birch Park	Birch Place	Greenwood	
Bracken Park	Bracken Court	Duncraig	
Bramston Park	Bramston Vista	Burns Beach	
Burns Park	Ocean Parade	Burns Beach	
Byrne Park	Byrne Close	Padbury	
Calectasia Park	Calectasia Street	Greenwood	
Captain Park	Captain Court	Heathridge	
Carr Park	Carr Crescent	Warwick	
Cliff Park	Cliff Street	Marmion	

Name	Address	Suburb
Clifford Coleman Park	Troy Avenue	Marmion
Cliverton Park	Cliverton Court	Marmion
Colac Park	Colac Way	Duncraig
Cornish Park	Cornish Avenue	Woodvale
Craigie Park	Barradine Way	Craigie
Culwalla Park	Culwalla Close	Kallaroo
Cumberland Park	Cumberland Way	Beldon
Doveridge Park	Doveridge Drive	Duncraig
Ellison Park	Ellison Drive	Padbury
Filbert Park	Filbert Street	Greenwood
Fraser Park	Fraser Way	Padbury
Gerda Park	Gerda Court	Greenwood
Glenmere Park	Glenmere Road	Warwick
Glenunga Park	Glenunga Way	Craigie
Greenwich Park	Greenwich Court	Kingsley
Grenfell Park	Grenfell Avenue	Duncraig
Gull Park	Gull Street	Marmion
Hakea Park	Hakea Place	Sorrento
Harcourt Park	Harcourt Drive	Hillarys
Hartley Park	Hartley Court	Greenwood
Hawick Park	Hawker Avenue	Warwick
Hillwood Park	Hillwood Avenue	Warwick
Hughes Park	Hughes Court	Padbury
Huxley Park	Huxleys Trail	Burns Beach
Karuah Park	Karuah Way	Greenwood
Kelvin Park	Campion Street	Duncraig
Kingfisher Park	Kingfisher Way	Kingsley
Kurrajong Park	Kurrajong Place	Greenwood
Lakeview Park	Lakeview Drive	Edgewater
Lanark Park	Lanark Mews	Duncraig
Leaside Park	Leaside Way	Greenwood
Leichhardt Park No2	Leichhardt Avenue	Padbury
Macarthur Park	Macarthur Avenue	Padbury
Mackay Park	Mackay Way	Hillarys
Magpie Reserve	Ozone Road	Marmion
Mayflower Park	Mayflower Crescent	Hillarys
Mckinlay Park	Macarthur Avenue	Padbury
Mcrae Park	Mcrae Court	Padbury
Minchin Park	Minchin Court	Padbury
Monkhouse Park	Monkhouse Way	Hillarys
Monument Park	Monument Drive	Beldon
Mueller Park	Mueller Court	Padbury
Nalpa Park	Kenny Drive	Duncraig
Nekaya Park	Nekaya Way	Duncraig
New Cross Park	New Cross Road	Kingsley
Nimrod Park	Nimrod Place	Hillarys
Oakapple Park	Oakapple Drive	Duncraig
	Camberwarra Drive	Craigie
Otway Park		
Otway Park Paterson Park	Paterson Place	Padbury

Name	Address	Suburb
Plover Park	Plover Way	Kingsley
Ranford Park	Ranford Way	Hillarys
Readshaw Park	Readshaw Road	Duncraig
Reserve For Recreation	Grand Ocean Entrance	Burns Beach
Ridge Park	Ridge Close	Edgewater
Sanday Park	Hawker Avenue	Warwick
Santa Ana Park	Aberfoyle Heights	Currambine
Southport Park	Southport Loop	Burns Beach
Spoonbill Park	Spoonbill Grove	Kingsley
Stillwater Park	Reflection Close	Edgewater
Strathaven Park	Strathaven Crescent	Greenwood
Talbot Park	Talbot Drive	Kingsley
Trailwood Park	Trailwood Drive	Woodvale
Tuart Park	Tuart Trail	Edgewater
Village Park	Village Walk	Ocean Reef
Waltham Park	Waltham Street	Mullaloo
Wandina Park	Wandina Place	Duncraig
Warner Park	Warner Place	Greenwood
Wentworth Park	Wentworth Way	Padbury
Windich Park	Windich Court	Padbury
Windmill Park	Windmill Circle	Burns Beach
Woodland Park	Woodland Loop	Edgewater
Candlewood Park	Candlewood Boulevard	Joondalup
Carnaby Reserve	Medinah Mews	Connolly
Clare Park	Clare Street	Sorrento
Coastal Foreshore Reserve Marmion	West Coast Drive	Marmion
Culwalla Park	Culwalla Close	Kallaroo
Dardanus Park	Telstar Place	Heathridge
Gunida Park	Gunida Way	Mullaloo
Harman Park	Harman Road	Sorrento
Harman Park	Harman Road	Sorrento
Harman Park	Harman Road	Sorrento
Kilrenny Park	Gilmerton Way	Greenwood
Kuta Park	Kuta Circuit	Iluka
Ledge Park	Ledge Place	Sorrento
Lookout Park	Quarry Ramble	Edgewater
Madana Park	Madana Place	Craigie
Madana Park	Madana Place	Craigie
Mandalay Park	Mandalay Place	Craigie
Okely Park	Joondalup Drive	Edgewater
Pentland Park	Pentland Avenue	Duncraig
Sweeney Park	Sweeney Way	Padbury

Appendix 4



Standard Dieback Signage Protocol for Use May 2008



This document sets out the protocols for the use of Standard Phytophthora Dieback Signage on all land tenures in Western Australia.





T: 08 9400 4000 F: 08 9300 1383 Boas Avenue Joondalup WA 6027 PO Box 21 Joondalup WA 6919

joondalup.wa.gov.au This document is available in alternate formats upon request.